

A Functional Evolutionary Approach to Identify Determinants of Nucleosome Positioning: A Unifying Model for Establishing the Genome-wide Pattern

Amanda L. Hughes,^{1,3} Yi Jin,^{2,3} Oliver J. Rando,^{1,*} and Kevin Struhl^{2,*}

¹Department of Biochemistry and Molecular Pharmacology, University of Massachusetts Medical School, 364 Plantation Street, Worcester, MA 01605, USA

²Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, MA 02115, USA

³These authors contributed equally to this work

*Correspondence: oliver.rando@umassmed.edu (O.J.R.), kstruhl@hms.harvard.edu (K.S.)

<http://dx.doi.org/10.1016/j.molcel.2012.07.003>

SUMMARY

Although the genomic pattern of nucleosome positioning is broadly conserved, quantitative aspects vary over evolutionary timescales. We identify the *cis* and *trans* determinants of nucleosome positioning using a functional evolutionary approach involving *S. cerevisiae* strains containing large genomic regions from other yeast species. In a foreign species, nucleosome depletion at promoters is maintained over poly(dA:dT) tracts, whereas internucleosome spacing and all other aspects of nucleosome positioning tested are not. Interestingly, the locations of the +1 nucleosome and RNA start sites shift in concert. Strikingly, in a foreign species, nucleosome-depleted regions occur fortuitously in coding regions, and they often act as promoters that are associated with a positioned nucleosome array linked to the length of the transcription unit. We suggest a three-step model in which nucleosome remodelers, general transcription factors, and the transcriptional elongation machinery are primarily involved in generating the nucleosome positioning pattern *in vivo*.

INTRODUCTION

In living cells, nucleosome positions are influenced by intrinsic DNA sequence preferences due to the thermodynamic costs associated with wrapping stiff DNA around the histone octamer (Drew and Travers, 1985; Jiang and Pugh, 2009; Radman-Livaja and Rando, 2010). In addition, a wide variety of proteins can affect nucleosome positions and occupancy, most notably ATP-dependent chromatin-remodeling complexes. The relative importance of DNA sequence and protein factors in determining nucleosome positioning has been subject to considerable debate. *In vitro* reconstitution of genomic DNA into nucleosomes by salt dialysis recapitulates gross variation in nucleosome occupancy in yeast and in humans—AT-rich sequences such

as those found at yeast promoters are intrinsically nucleosome depleted (Kaplan et al., 2009; Sekinger et al., 2005; Zhang et al., 2009), whereas the GC-rich sequences prevalent at human promoters are intrinsically nucleosome enriched (Valouev et al., 2011). These studies typically find little role for intrinsic preferences in precise nucleosome positioning, although the enrichment of particular sequence features (10 bp periodicity of AA/AT/TA dinucleotides) at the +1 position in budding yeast has nonetheless led to forceful (Kaplan et al., 2009, 2010; Segal et al., 2006) but disputed (Fan et al., 2010; Stein et al., 2009; Weiner et al., 2010; Zhang et al., 2009, 2010) claims that intrinsic DNA sequence preferences play a major determining role in nucleosome positioning.

Conversely, several experimental approaches, largely in budding yeast, have revealed a key role for proteins in establishing nucleosome positions *in vivo*. While *in vitro* reconstitution of DNA into nucleosomes does not properly establish nucleosome positions at *PHO5*, addition of yeast whole-cell extract enables more accurate assembly of nucleosomes at this locus (Korber and Horz, 2004). Genome-wide analysis subsequently showed that one or more ATP-dependent activities in yeast whole-cell extract can assemble nucleosomes in positions that resemble, but do not completely coincide with, *in vivo* positioning (Zhang et al., 2011), thereby demonstrating a major role for nucleosome-remodeling complexes in nucleosome positioning. Decades of biochemical studies have identified many specific proteins and protein complexes capable of altering nucleosome positions on DNA *in vitro* (Clapier and Cairns, 2009), and increasingly these factors are being implicated in proper nucleosome positioning *in vivo* (Gkikopoulos et al., 2011; Whitehouse et al., 2007; Whitehouse and Tsukiyama, 2006). For example, the ATP-dependent remodeling enzymes Chd1, Isw1, and Isw2 globally affect nucleosome positioning *in vivo*, as their deletion in yeast leads to nearly complete loss of nucleosome positioning downstream of the +2 nucleosome of coding regions (Gkikopoulos et al., 2011).

In a genetic approach to this problem, diploid hybrids between the closely related species *S. cerevisiae* and *S. paradoxus* have been used to determine to what extent divergent nucleosome positioning on specific orthologous genes can be attributed to *cis* or *trans* factors, with the majority of chromatin changes between these species being attributed to poly(dA:dT) elements




Figure 1. Functional Evolutionary Dissection of Chromatin-Establishment Mechanisms

(A) Schematic of experimental design. Yeast artificial chromosomes are constructed carrying sequence from species such as *K. lactis*, and introduced into *S. cerevisiae*. Comparison of nucleosome-mapping data between the same sequence in two different environments (its endogenous genome, and in *S. cerevisiae*) can be used to disentangle DNA-driven from *trans*-mediated aspects of chromatin organization.

(B) Chromosomal complement of parental *S. cerevisiae* (AB1380) and three different YAC-bearing strains, as indicated. Pulsed-field gel electrophoresis of YAC-bearing strains, as indicated.

(C and D) Examples of nucleosome-mapping data from two genes. Blue line indicates nucleosome-mapping data from wild-type *K. lactis* (Tsankov et al., 2010), red line shows data from the same sequence carried on a YAC in *S. cerevisiae*.

(E and F) Data for all *K. lactis* genes on all three YACs. (E) shows data for all genes from wild-type *K. lactis*, with genes sorted by NDR width, while (F) shows data from these genes on YACs, sorted identically. Black indicates no sequencing reads, yellow intensity indicates number of sequencing reads. C and D indicate the example genes shown above.

the nuclear environment to nucleosome positioning *in vivo*. This approach relies on the finding that there are species-specific differences in parameters of nucleosome positioning in a variety of yeast species, even though the general pattern is highly conserved (Tsankov et al., 2010). Specifically, we compare nucleosome maps of artificial chromosomes (YACs) containing large, heterologous genomic regions from different yeast species in *S. cerevisiae* with maps of the same regions in their native organism (Figure 1A). In principle, features that change in the context of *S. cerevisiae* are determined by protein factors that are functionally distinct in the two species, whereas features that are retained when the foreign yeast DNA is present in *S. cerevisiae* are due either to intrinsic DNA sequence or to

conserved *trans*-acting regulators. For example, when the *S. cerevisiae* *HIS3-PET56* region is introduced into *S. pombe*, it retains the nucleosome-depleted promoter region, but not the positions of nucleosomes in the coding region (Sekinger et al., 2005). In addition, the generation of fortuitous functional elements arising from heterologous genomic sequences makes it possible to address mechanistic issues that are presumably free of evolutionary constraints. Here, we show that nucleosome spacing is established *in trans*, and that promoter nucleosome

at promoters (Tirosh et al., 2010). However, *S. cerevisiae* and *S. paradoxus* differ very little in bulk aspects of chromatin architecture. In contrast, chromatin structure exhibits far greater differences between more divergent species: for example, average nucleosome spacing differs by ~15–20 bp between *S. cerevisiae* and *K. lactis* (last common ancestor ~150 million years ago) (Heus et al., 1993; Tsankov et al., 2010).

Here, we describe a functional evolutionary approach to systematically dissect the contributions of DNA sequence and

depletion can be established either by intrinsic sequence cues or by *trans*-acting factors. Further, we find that +1 nucleosome positioning is most likely established by some aspect of the transcriptional machinery, and positioning of more downstream nucleosomes in the mRNA coding region is linked to Pol II elongation. Based on results presented here and elsewhere, we propose a unifying, three-step model for how nucleosome positions are established in vivo.

RESULTS

Generation of *S. cerevisiae* Strains Harboring Artificial Chromosomes with Large Segments of Foreign Yeast DNA

To generate yeast artificial chromosomes (YACs, Figure 1A), genomic DNA from *K. lactis*, *K. waltii*, and *D. hansenii* was sheared to ~100–200 kb average size and ligated to the pYAC4 vector carrying sequences for *S. cerevisiae* telomeres, centromere, and origin of replication, as well as two selectable markers. YACs were transformed into wild-type *S. cerevisiae* and confirmed by pulsed-field gel electrophoresis (Figure 1B). Furthermore, both ends of YACs containing foreign yeast DNA inserts were validated by DNA sequencing. In total, we generated seven strains carrying distinct YACs from the three species, with an average insert length of ~140 kb (see Table S1 online). YAC strains were grown in conditions identical to those previously used for mapping nucleosomes in these four species (Tsankov et al., 2010), and formaldehyde crosslinked chromatin was digested to ~80% mononucleosomes using micrococcal nuclease (Yuan et al., 2005). Mononucleosomal DNA was analyzed by deep sequencing as previously described (Shivaswamy et al., 2008; Tsankov et al., 2010; Weiner et al., 2010). Figures 1C and 1D show nucleosome-mapping data for two genes from *K. lactis*, with data from wild-type *K. lactis* in blue (“endogenous”) and data for these same genes in the context of a YAC-carrying *S. cerevisiae* strain in red (“YAC”). Notable in these views are a number of well-described aspects of fungal chromatin structure—in the endogenous context, nucleosomes are generally well positioned (nucleosome peaks are well separated and exhibit high peak to trough ratios), and both genes have a nucleosome-depleted region (NDR) that contains the gene’s promoter.

Promoter NDRs Are Largely Maintained in a Foreign Species in a Manner Strongly Correlated with Poly(dA:dT) Tracts

The endogenous positions of promoter NDRs were largely maintained in the YACs (Figures 1E and 1F)—50% and 56% of *D. hansenii* and *K. lactis* NDRs, respectively, were located within 50 bp of their endogenous position, and for both species’ sets of YACs only 13% of NDRs did not overlap the endogenous NDR at all. Furthermore, the extent of the NDR, which varies considerably among genes, is largely maintained in the YAC-containing strains. These data are consistent with the view that nucleosome depletion at fungal promoters is largely programmed by genomic sequence. However, the average extent of depletion over promoters is not as great in YACs as in wild-type (Figures 1C–1F, and Figures S1A and

S1B), potentially as a consequence of the reduced expression of YAC genes (see below). This observation suggests that some of the depletion at promoters is not intrinsically determined by DNA sequence, consistent with the previous observation that nucleosome depletion at promoters is more pronounced in vivo than in vitro (Kaplan et al., 2009; Zhang et al., 2009, 2011).

To further investigate the role for intrinsic sequence cues in establishing nucleosome depletion, we sorted genes by the difference in nucleosome occupancy over the proximal NDR between endogenous genes and YACs (Figure 2A, Figures S1C and S1D). Notably, in both *K. lactis* and *D. hansenii* we observed very few genes with lower nucleosome occupancy in the YAC context, with the majority of promoters showing a range from little change to substantially increased nucleosome occupancy. Genes that maintained the same level of nucleosome depletion in the YAC as in wild-type were characterized by promoter sequences with greater numbers of long poly(dA:dT) elements (Figures 2B and 2C, see the Experimental Procedures), consistent with the idea that these sequences intrinsically program nucleosome depletion in any context. Genes from *D. hansenii* tended to exhibit much less dramatic nucleosome depletion at promoters in the YAC context than in their endogenous context (Figure S1B). This is consistent with our prior observation (Tsankov et al., 2010, 2011) that *D. hansenii* promoters have fewer poly(dA:dT) sequences than most other organisms in the *Hemiascomycota* phylogeny (potentially due to their ecological niche in high-salt environments) and with the hypothesis that *D. hansenii* promoters are more often established by *trans*-acting proteins such as General Regulatory Factors (GRFs). In this regard, promoters that gained substantial nucleosome occupancy when carried in the YAC sometimes, but not always, contained known binding motifs for transcription factors we previously (Tsankov et al., 2010, 2011) inferred to be GRFs in *D. hansenii*, but not in *S. cerevisiae* (Figure 2D). Together, these results indicate that intrinsic sequence determinants (or conserved *trans*-acting factors) play a major role in generating nucleosome depletion at fungal promoters, and that poly(dA:dT) tracts are the primary DNA sequence determinant.

Nucleosome Positions Differ Markedly in the Endogenous and YAC-Containing Strains

In contrast to the widespread but not universal conservation of NDRs, a given DNA sequence is generally packaged differently when carried in the endogenous species or in *S. cerevisiae*. Nucleosome positions change markedly in the YAC strains—the +1 nucleosome can be found near to (Figure 1C), upstream (see below), or downstream (Figure 1D) of its endogenous location, while nucleosomes farther downstream of the +1 occur farther and farther away from their endogenous locations. By definition, differences in nucleosome positioning of a given genomic region in the endogenous organism or in *S. cerevisiae* cannot be due to intrinsic DNA sequence, but rather *trans*-acting factor(s). These measured differences are not secondary to technical artifacts such as differences in MNase digestion, as we observe remarkably consistent results for *S. cerevisiae* genes for the various YAC data sets (Figure S2). Interestingly, the




Figure 2. Promoter Nucleosome Depletion Is Maintained over Poly(dA:dT) Elements

(A) *D. hansenii* genes sorted by the extent of change in nucleosome occupancy over the NDR. Left panel shows differences in nucleosome occupancy between *D. hansenii* and YACs for 114 genes—blue indicates increased nucleosome occupancy in the YAC relative to endogenous context. Middle and right panels show nucleosome-mapping data for endogenous *D. hansenii* sequences and for YACs, as indicated.

(B) Strength of poly(dA:dT) element (Field et al., 2008; Tsankov et al., 2010) for genes, ordered as in (A). Forty gene running window average is shown.

(C) An example of a gene with little change in nucleosome depletion between endogenous and YAC contexts. Sequence from this stable NDR contains multiple poly(dA:dT) elements, as indicated in red.

(D) An example of a gene exhibiting dramatically increased nucleosome occupancy at the native NDR when carried on YAC. Here, this NDR includes few poly(A) elements, and carries a binding site for Cbf1, which has nucleosome-evicting activity in *D. hansenii* but not in *S. cerevisiae* (Tsankov et al., 2010, 2011).

average deviation in chromatin structure between genes in their endogenous context and in the YAC was quite different for the three species studied—*K. lactis* genes appeared closest to their native structure when in YACs, whereas chromatin structure of *K. waltii* sequences in YACs appeared random with respect to genic structure (data not shown).

Nucleosome Spacing Is Determined by Protein Factors in the Host Organism, Not DNA Sequence

As observed by MNase cleavage of bulk chromatin, most nucleosomes in any given species are found in arrays in which the linker regions between adjacent nucleosomes are similar in size. Interestingly, nucleosome spacing is substantially different




Figure 3. Nucleosome Spacing Is Set In trans

(A) Averaged data for all *K. lactis* genes on YACs 1–3. Genes are aligned by the +1 nucleosome position as defined in Tsankov et al. (2010), and data from either wild-type *K. lactis* or from the YAC strains are averaged for 184 genes, as indicated.

(B) *K. lactis* sequences adopt *S. cerevisiae* spacing when carried in *S. cerevisiae*. Nucleosome positions were called, and the distribution of all internucleosomal distances (center to center) is shown for 184 *K. lactis* genes from wild-type or in the YACs. Similar distributions for *S. cerevisiae* nucleosome positioning from wild-type and YAC-containing strains indicate that YACs do not perturb host chromatin state (see also Figure S2).

between *S. cerevisiae*, with an average internucleosomal distance of ~165 bp, and *K. lactis*, with an average spacing of ~178 bp (Heus et al., 1993; Tsankov et al., 2010). We therefore used our YAC data set to assess whether nucleosome spacing over *K. lactis* genes is established by DNA sequence, or by the nuclear environment. As can be appreciated in Figures 1C and 1D, nucleosome spacing appears shorter over *K. lactis* genes when carried in *S. cerevisiae*, relative to the endogenous spacing. Figure 3A shows the average nucleosome data for all *K. lactis* genes present on one of the 3 YACs, aligned by the endogenous location of the +1 nucleosome. Average nucleosome spacing decreases when these genes are carried in YACs. The distribution underlying this average trend is quantified in Figure 3B. Here, we called nucleosome positions (Weiner et al., 2010), then plotted the distribution of all internucleosomal distances as indicated. *K. lactis* genomic sequence in its endogenous context is packaged with nucleosomes occurring every 178 bp, whereas the same sequence in the *S. cerevisiae* *trans* environment exhibits ~165 bp nucleosome spacing, precisely the same spacing observed over native *S. cerevisiae* genes. Importantly, we observed no change in the spacing of *S. cerevisiae* genes between wild-type yeast and our YAC strains, indicating no artifactual effects on nucleosome spacing from, for example, MNase titration level (Figure 3B, Figure S2). The primary role for proteins in determining internucleosomal spacing is not surprising, as different cell types in multicellular organisms (sharing identical genomes) can exhibit different nucleosome spacing (Van Holde, 1989). Importantly, the observation that internucleosomal spacing depends on protein factors

means that the precise positions for the vast majority of nucleosomes are not determined by intrinsic DNA sequence.

The Position of the +1 Nucleosome Is Not Determined by DNA Sequence but Rather Is Mechanistically Linked to Transcriptional Initiation

The claim that the +1 nucleosome is positioned by DNA sequence (Segal et al., 2006) has been subject to debate, not least because in vitro reconstitution experiments reveal no significant recovery of +1 nucleosome positioning. Alternatively, it has been proposed that the +1 nucleosome is positioned by either transcription factors such as Rap1, Abf1, and Reb1 (Kornberg and Stryer, 1988; Zhang et al., 2009, 2011); NDRs (Mavrich et al., 2008; Yuan et al., 2005); or the preinitiation complex (Zhang et al., 2009).

Interestingly, although the average position of the +1 nucleosome is similar between YACs and the endogenous context for both *K. lactis* and *D. hansenii* genes (see Figure 3A, Figure S1), examination of individual genes shows that +1 positioning is highly variable for the same sequence in two different nuclear environments. The distribution of +1 nucleosome shifts for genes carried on YACs was far more variable than the experimental variability measured using the background of *S. cerevisiae* genes—while only 17% (14%–21% in different YAC strains) of *S. cerevisiae* +1 nucleosomes appeared >20 bp apart between strains, 45% of *K. lactis* and 57% of *D. hansenii* +1 nucleosomes in the YAC strains shifted at least 20 bp from their endogenous location (Figure S3). +1 nucleosomes could shift in either direction on YACs (Figure 4, Figure S3), although in *K. lactis* YACs




Figure 4. +1 Nucleosome Shifts Associated with Transcriptional Changes

(A and B) Nucleosome data and RNA-Seq data are shown for *K. lactis* and *D. hansenii* genes in wild-type and YACs, as indicated. RNA-Seq data for YAC-derived transcripts are normalized independently from *S. cerevisiae* transcripts here—see Figures S4B and S4C for data normalized genome-wide.

(C–E) Examples of +1 nucleosome shifts associated with changes in transcription. (C) shows a moderate upstream shift in a +1 nucleosome with a similar change in transcript length, while (D) and (E) show large-scale NDR gain/loss with associated changes in transcription. Schematic interpretation of the nucleosome positioning for the endogenous gene is shown in blue above the rectangle, nucleosome positioning in the YAC is shown in red below the rectangle. Arrows indicate inferred TSSs (note that RNA-sequencing data are not strand specific, but TFIIIB-mapping data support our inferred TSSs)—the furthest 5' RNA in (E), for example, derives from the upstream gene as opposed to a divergent promoter.

these shifts were biased toward upstream shifts (Figure 3A). Thus, our observations demonstrate that pronucleosomal sequences do not “program” the position of the +1 nucleosome in vivo.

The strong correspondence between +1 nucleosome positioning and transcriptional start sites in many species (Jiang and Pugh, 2009) led us to consider the hypothesis that changes in transcriptional activity might underlie the repositioning of the +1 nucleosomes (Zhang et al., 2009). We therefore carried out deep sequencing of RNA isolated from *D. hansenii*, *K. lactis*, and the *S. cerevisiae* YAC strains in this study, and carried out ChIP-Seq for TFIIIB localization in the YAC-containing strains (a full analysis of these data will be published separately). Alignment of RNA-Seq data from wild-type strains with nucleosome-mapping data confirmed prior predictions that the positioning of +1 nucleosomes with respect to a gene’s transcription start site (TSS) varies between these species (Tirosh et al., 2007; Tsankov et al., 2010)—transcription begins further inside the +1 nucleosome in *K. lactis* than in *D. hansenii* (Figure S4A).

Comparing endogenous to YAC-based gene expression, we found on average that genes on YACs were expressed at lower

levels than host genes—average sequencing reads per kilobase of coding sequence for YACs were ~40% of the average value for endogenous RNAs—consistent with extensive promoter sequence divergence between species resulting in widespread misinterpretation of exogenous regulatory information by the *S. cerevisiae* transcriptional machinery (Figures S4B, S4C, and S4E). In general, we found a good correlation between expression levels for genes in their endogenous genome versus expression from the YACs (Figure S4D)—genes expressed at high levels in *K. lactis* remained the most highly expressed genes when carried on YACs, but were expressed at lower levels relative to *S. cerevisiae* genes. In *D. hansenii*, we also observed increased expression of intergenic regions in the YACs (Figure S4C and see below), again indicating evolutionary divergence in transcriptional control sequences (e.g., loss of transcriptional termination signals and/or gain of cryptic promoters).

Consistent with a relationship between +1 nucleosome positioning and TSSs, we found that the 5' ends of RNAs in YACs shifted on average toward a *S. cerevisiae*-like location relative to the +1 nucleosome (Figures 4A and 4B)—*K. lactis* RNAs started farther upstream in the YAC, whereas *D. hansenii* RNAs

Molecular Cell

Unifying Model for Nucleosome Positioning

started farther downstream. Furthermore, +1 nucleosome shifts largely were accompanied by coherent shifts in inferred TSSs. These include ~90 examples such as that seen in Figure 4C, in which both RNA-Seq data and the +1 nucleosome for a given gene shift in the same direction. This is visualized in Figure S5A—RNA-Seq data for *K. lactis* genes exhibiting no +1 nucleosome shift, and for those exhibiting upstream +1 nucleosome shifts, are averaged for both endogenous RNA expression and YAC-based expression. Despite no average difference in 5' ends of transcripts between these two classes in the endogenous case, we find that genes exhibiting upstream shifts in the +1 nucleosome also showed more strongly 5'-shifted transcripts relative to genes without a +1 nucleosome shift, consistent with the idea that there is a mechanistic coupling between +1 nucleosome positioning and transcriptional initiation. Furthermore, we also used 5' RACE to more precisely map TSSs for four *K. lactis* genes in their endogenous context and in the YAC, confirming that +1 nucleosome shifts corresponded to shifts in the location of the TSS (Figures S5B–S5E). These observations provide functional evidence for a mechanistic linkage between nucleosome positioning and transcriptional initiation, although they do not establish the cause-and-effect relationship between these two processes.

Generation of NDRs in Foreign Coding Regions via Fortuitous Interactions of *S. cerevisiae* Activator Proteins

More dramatic cases of nonconserved nucleosome positioning are observed, particularly in *D. hansenii*-derived YACs, in which many NDRs arise in coding regions (Figures 4D and 4E, Figures 5A and 5B, Figures S6A and S6B). The sequences underlying these NDRs are not associated with poly(dA:dT) elements (data not shown), as expected given that they do not intrinsically form NDRs in their host genomic context (Figure 5B, blue line). Interestingly, these NDRs are associated with TFIIB binding (Figures 5C and 5D) and concomitant changes in RNA abundance (Figures 4D and 4E, Figure 5A, Figure S6A), indicating a wholesale functional change in which a coding sequence from one species (*D. hansenii*) is used as a promoter in a foreign species (*S. cerevisiae*). These NDRs are most likely determined by *S. cerevisiae* transcription factors that fortuitously recognize and functionally act on foreign DNA sequences that do not act as promoters in the native organism. In other words, DNA-binding transcriptional activator proteins recruit nucleosome-remodeling complexes to these fortuitously recognized sites, thereby evicting histones and generating an NDR. These NDRs are associated with varying levels of TFIIB and RNA transcripts, presumably depending on the quality of TATA elements and other core promoter sequences in the vicinity.

Fortuitous Coding Region NDRs Are Associated with Typical Nucleosome Patterns

The existence of fortuitous and presumably evolutionarily meaningless promoters in *D. hansenii* coding regions in the context of *S. cerevisiae* cells makes it possible to determine the role of transcription in establishing the nucleosome positioning pattern. Strikingly, these coding region NDRs are asso-

ciated with a typical nucleosome pattern of highly positioned +1 and -1 nucleosomes as well as progressively less positioned downstream nucleosomes (Figure 5B). Thus, in the absence of any intrinsic nucleosome-destabilizing sequences, transcription factors and associated cofactors are sufficient to generate a nucleosome positioning pattern that is very similar to the standard pattern at endogenous promoters. Furthermore, at such fortuitous NDRs, the extent of the positioned array is linked to the length of the RNA transcript (Figure 5A, Figure S6C), strongly suggesting a role for transcriptional elongation in the generation of the nucleosomal pattern. These results demonstrating a functional role for transcription-related events appear to conflict with the conclusion that nucleosome-remodeling complexes are sufficient to establish aspects of the nucleosome positioning pattern in the absence of transcription (Zhang et al., 2011). However, these observations are not mutually exclusive, and indeed are complementary, as both mechanisms are likely to contribute to establishing the nucleosome pattern.

DISCUSSION

A Functional Evolutionary Approach to Address the Determinants of Molecular Phenomena In Vivo

Here, we used a functional evolutionary approach to systematically dissect the role for *cis*-acting sequence elements and *trans*-acting proteins in establishment of nucleosome positioning in fungi. This approach, which is based on species-specific differences in parameters of nucleosome positioning in a variety of yeast species (Tsankov et al., 2010), involves placing large segments of foreign yeast DNA in *S. cerevisiae* and comparing molecular properties in such strains with those in the native organism. In principle, nonconserved properties are determined by protein factors that are functionally distinct in the two species, whereas conserved properties are due either to DNA sequence or to conserved *trans*-acting regulators. The use of yeast artificial chromosomes to carry the foreign yeast DNA makes it possible to examine many genes at once, and hence to obtain information that is both statistically robust and permits one to identify many examples of new phenomena. Furthermore, the ability to generate fortuitous functional events (e.g., the NDRs in *D. hansenii* coding regions) that do not occur in the native organisms makes it possible to address mechanistic questions in a manner that is, most likely, independent of evolutionary history. An extension of this approach should also permit one to identify factors responsible for the species-specific behavior, specifically by replacing a candidate factor by its homolog in the foreign species and examining whether the pattern resembles that of the foreign species.

More generally, this functional evolutionary approach should allow for elucidating the determinants of other molecular phenomena that are broadly conserved but show species-specific differences. For example, more detailed analysis of the RNA generated in the YAC-containing strains with the corresponding endogenous yeast species should be informative of determinants of 5' and 3' end formation, splicing, and half-lives. As such, this approach combines the virtues of evolutionary comparison and classic functional genetic analysis.




Figure 5. Characterization of Fortuitous NDRs in YACs

(A) Example of a fortuitous NDR that occurs only in the YAC but not in the native genome, and is associated with transcription. This fortuitous NDR occurs in the middle of a *D. hansenii* coding region and is associated with two shorter, divergent transcripts in the YAC context (data cover 2.2 kb of sequence). Note that nucleosome organization correlates with transcript length—rightmost transcript shows greater nucleosome positioning at the 5' end than at the 3' end of the transcript.

(B) Fortuitous NDRs are generally associated with well-positioned ± 1 nucleosomes. Averaged data for 120 NDRs observed in *D. hansenii* YACs but not in the endogenous context, as indicated.

(C and D) Fortuitous NDRs represent functional promoters. (C) shows TFIIIB ChIP-Seq data from YAC-bearing strain for the genomic locus shown in (A), while (D) shows averaged data for all fortuitous NDRs. Note that TFIIIB localization in the endogenous context could not be obtained, as our anti-TFIIIB antibody does not recognize TFIIIB from *D. hansenii*.

A Three-Step Model for Establishing the Nucleosome Positioning Pattern In Vivo


Based on results presented here and elsewhere, we propose a three-step model (Figure 6) for how nucleosome positioning is established in eukaryotic organisms. The first step involves the generation of an NDR, which can occur either by transcription factors and their recruited nucleosome-remodeling complexes and/or by poly(dA:dT) sequences that intrinsically disfavor nucleosome formation. Even at poly(dA:dT)-containing promoters, it is likely that transcriptional machinery contributes to nucleosome depletion, as nucleosome depletion is more pronounced in vivo than in vitro (Kaplan et al., 2009; Zhang et al., 2009), and nucleosome-remodeling complexes enhance the depletion in vitro (Zhang et al., 2011). In this sense, intrinsic

programming of NDRs represents a specialized mechanism that is used frequently (*S. cerevisiae*), moderately (*D. hansenii*), or rarely (*D. melanogaster*), depending on the species.

In the second step, nucleosome-remodeling complexes recognize the NDRs and generate highly positioned nucleosomes flanking the NDR. Strong positioning could, in principle, arise simply from the boundary of the NDR and/or from sequence preferences of the nucleosome remodelers. Indeed, it has been argued that this step does not require transcription factors or transcription per se (Zhang et al., 2011), although it is important to note that there is overall poor correspondence between +1 nucleosome positioning observed in vivo and that recapitulated using ATP-dependent extracts in the absence of transcription (Zhang et al., 2011) (Figure S7). In this regard,


Step 1: NDR generation

(intrinsic, and/or activator-dependent)



Step 2: Positioning nucleosomes flanking the NDR

(established by nucleosome remodelers, likely fine-tuned by PIC)



Step 3: Transcriptional elongation mediated nucleosome positioning

(elongating RNAPII, Chd1, Isw1, etc.)




Figure 6. Three-Step Model for Establishment of Nucleosome Positioning In Vivo

A unifying three-step model for how nucleosome positioning pattern is generated in eukaryotic organisms. The first step is the generation of an NDR, either by poly(dA:dT) elements and/or by transcription factors and their recruited nucleosome remodeling complexes. In the second step, nucleosome remodeling complexes recognize the NDRs and generate highly positioned nucleosomes flanking the NDR; and the RNA polymerase II preinitiation complex fine-tunes the position of the +1 nucleosome. In the final step, positioning of the more downstream nucleosomes depends on transcriptional elongation and the recruitment of nucleosome remodeling activities and histone chaperones by the elongating RNA polymerase II machinery.

Zhang et al. compared nucleosome positioning generated by ATP-dependent extracts with the nucleosome positions measured from yeast lysed without crosslinking and allowed to redistribute prior to crosslinking. Indeed, we find mediocre correspondence between the “native” nucleosome positions from Zhang et al. and true in vivo nucleosome positions generated from crosslinked yeast (Figure S7), so the ability of whole-cell extracts to recover these “native” positions in the absence of transcription does not have any bearing on the question of whether in vivo positioning is influenced by transcription prior to lysis of cells. Although nucleosome remodelers can generate somewhat positioned nucleosomes flanking the NDR and unquestionably perform far better than salt dialysis, they apparently are insufficient to generate the precise in vivo nucleosome positions, particularly for the +1 nucleosome (Figure S7).

Here, the strong, and species-specific, spacing relationship between the +1 nucleosome and mRNA start site that is observed both in the native and YAC strains indicates that there is a mechanistic connection between transcriptional initiation and the location of the +1 nucleosome. Given the strong in vivo positioning of both the preinitiation complex and the +1 nucleosome, a spacing relationship between these two entities requires that at least one of these is anchored to a specific location,

thereby permitting a defined location for the second entity. As discussed above, nucleosome remodeling complexes alone are insufficient to generate proper positioning of the +1 nucleosome, and hence sequence and nucleosome remodelers are insufficient to provide an anchor. In contrast, preinitiation complexes bound at core promoters are clearly sufficient to provide an anchor, with the location of the TBP bound to the TATA element or TATA-related sequence being the major determinant of the anchor point. From these considerations, and our finding that the TSS to +1 distance in YACs shifts to the *S. cerevisiae* spacing (Figure 5 and Figure S5), we suggest that the preinitiation complex plays a role in fine-tuning the position of the +1 nucleosome.

In the third step, positioning of downstream nucleosomes, with progressively less positioned nucleosomes downstream within the gene, depends on transcriptional elongation, and hence recruitment of nucleosome remodeling activities and histone chaperones by the elongating RNA polymerase II machinery. This elongation-dependent step explains why nucleosome remodeling complexes, though capable of weakly positioning nucleosomes flanking the NDR, are unable to position more downstream nucleosomes (Zhang et al., 2011). Conversely, yeast mutant strains lacking nucleosome remodeling complexes (Chd1 and Isw1) that are recruited to coding regions by elongating RNA polymerase show drastically reduced positioning of downstream nucleosomes but relatively normal positioning of the +1 and +2 nucleosomes (Gkikopoulos et al., 2011). Finally, a transcription-based step nicely helps to explain why nucleosome arrays occur largely in the transcribed direction even though highly positioned nucleosomes can occur both at the +1 and -1 position, as well as the curious observation that the decay of nucleosome positioning toward the center of genes displays a 5'/3' asymmetry (Vaillant et al., 2010); both of these observations are inconsistent with a pure packing-based model.

The above model can explain why the general pattern of nucleosome positioning is highly conserved among eukaryotes yet shows species-specific differences in various aspects of chromatin structure. These species-specific differences reflect the relative utilization of poly(dA:dT) sequences and hence intrinsic histone-DNA interactions, as well as differences in the enzymatic and recruitment properties of the nucleosome remodelers.

EXPERIMENTAL PROCEDURES

Growth Conditions

All cultures were grown in medium containing the following: SC -tryptophan -uracil (Sunrise Sciences) (0.2%), yeast extract (1.5%), peptone (1%), dextrose (2%), and adenine (0.01%), as previously described (Tsankov et al., 2010).

Preparation of YACs

Yeast chromosomal DNA was prepared in InCert agarose blocks (LONZA), with a final cell concentration of 2×10^9 cells/ml. Agarose blocks with intact chromosomal DNA were subjected to EcoRI partial digestion with a titrated Mg^{2+} concentration, followed by size fractionation using pulsed field gel electrophoresis (PFGE). Partially digested DNA fragments (~100–200 kb) were excised from the gel. YAC vector pYAC4 was purified by successive CsCl gradient ultracentrifugation and digested with BamHI and EcoRI, followed by calf intestine alkaline phosphatase treatment. Digested pYAC4 and partially digested yeast chromosomal fragments were ligated by T4

DNA ligase in agarose blocks. Prior to YAC transformation, ligated DNA was size fractionated again by PFGE, and DNA larger than 100 kb was excised from the gel. The excised gel slice was further digested with β -agarase, and ligated DNA was transformed into *S. cerevisiae* host cells (AB1380), using either spheroplast transformation protocol or standard yeast LiCl transformation method. Further details of YAC construction are available upon request.

Validation of YAC-Bearing Strains

Transformants with red color, which survived double selection (Ura⁺/Trp⁺) on AHC plates, were collected for validation. Chromosomal DNA of candidate strains was prepared in agarose blocks and resolved by PFGE (Figure 1B). Strains with desired YAC bands were selected, and terminal sequences from selected YAC clones were isolated and confirmed with DNA sequencing analysis (Riley et al., 1990).

Nucleosome Isolation and Illumina Deep Sequencing

Micrococcal nuclease (MNase) digestions were performed as previously described (Yuan et al., 2005). Briefly, 450 mL cultures were grown to OD₆₀₀ of ~0.5 at 30°C, 220 rpm. Cultures were fixed for 30 min at 30°C with 1.85% formaldehyde, then spheroplasted with 10 mg zymolyase (Cape Cod Associates) for 45 min at 30°C. Spheroplasts were subjected to 20 min of MNase digestion, and DNA was purified. MNase titrations were selected to obtain largely mononucleosomal DNA with some di- and trinucleosomal DNA apparent. Mononucleosomal DNA was gel purified (BioRad Freeze N' Squeeze) and used to create a library for deep sequencing on the Solexa 1G Genome Analyzer, as previously performed (Shivaswamy et al., 2008; Tsankov et al., 2010; Weiner et al., 2010). Briefly, DNA was blunt ended, A tailed, and ligated to Illumina genomic adapters, followed by a final PCR with a size-selecting gel purification.

Data Normalization and Nucleosome TSS Alignments

Reads from deep sequencing were mapped back to the relevant hybrid genome (*S. cerevisiae* plus the relevant species' chromosome) using blat. Uniquely mapping reads that had fewer than three mismatches were kept for analysis. Reads were extended by the crosscorrelations between those from the Watson and Crick strands, to create nucleosome peaks. Read count numbers were normalized to one by dividing each base read count by the genome-wide average read count per base. Gene alignments were carried out using the endogenous boundary of the +1 nucleosome (Tsankov et al., 2010). RNA-Seq data were treated similarly but without extending reads. RNA abundance for YAC-based transcripts was, on average, ~30%–40% (in reads per kb per million reads) of the RNA abundance of endogenous *S. cerevisiae* transcripts.

Nucleosome Calls

Template filtering (Weiner et al., 2010) was used to call the locations of nucleosomes.

5' RACE

Trizol (Invitrogen)-extracted RNA was enriched for mRNA on polyT magnetic beads (NEB). Calf intestinal phosphatase removed all phosphates prior to the hydrolysis of the mRNA cap to a phosphate with Tobacco Acid Pyrophosphatase (Epicenter). An oligo was ligated to the 5' end of the mRNA (T4 RNA Ligase), and the RNA was reverse transcribed (SuperScript III Reverse Transcriptase, Invitrogen) with a tailed random hexamer. The cDNA was amplified with a low-cycle PCR (Phusion, NEB) using primers matching the sequences added in the ligation and reverse transcription. A gene-specific PCR amplified the TSS sequence, which was cloned (StrataClone, Agilent) and sequenced.

ACCESSION NUMBERS

The Gene Expression Omnibus accession number for the data sets reported in this paper is GSE39011.

SUPPLEMENTAL INFORMATION

Supplemental Information includes seven figures and one table and can be found with this article at <http://dx.doi.org/10.1016/j.molcel.2012.07.003>.

ACKNOWLEDGMENTS

This paper is dedicated to the memory of Jonathan Widom, our friend and colleague who was passionately interested in nucleosome positioning and many other aspects of chromatin structure, and enjoyed vigorous scientific debate. His experimental and conceptual contributions were a major influence on this work, and we will miss our numerous memorable and enjoyable discussions. We thank Yajie Niu for assistance in CsCl gradient ultracentrifugation, Caitlin Reavey and Matthew Pipkin for helpful discussions on pulsed-field gel electrophoresis, and members of the Rando and Struhl labs for helpful discussions and comments. O.J.R. is funded by NIH grant GM079205, and K.S. is funded by GM030186. O.J.R. and K.S. conceived the experiments, and Y.J. and A.H. designed the experiments. Y.J. generated YACs and YAC-bearing strains, and A.H. carried out nucleosome isolation and deep sequencing, and 5' RACE. Y.J. carried out all RNA sequencing and TFIIB ChIP sequencing. A.H. and Y.J. carried out data analysis (with assistance from O.J.R. and K.S.). A.H., Y.J., O.J.R., and K.S. wrote the manuscript.

Received: February 16, 2012

Revised: June 11, 2012

Accepted: July 6, 2012

Published online: August 9, 2012

REFERENCES

- Clapier, C.R., and Cairns, B.R. (2009). The biology of chromatin remodeling complexes. *Annu. Rev. Biochem.* 78, 273–304.
- Drew, H.R., and Travers, A.A. (1985). DNA bending and its relation to nucleosome positioning. *J. Mol. Biol.* 186, 773–790.
- Fan, X., Moqtaderi, Z., Jin, Y., Zhang, Y., Liu, X.S., and Struhl, K. (2010). Nucleosome depletion at yeast terminators is not intrinsic and can occur by a transcriptional mechanism linked to 3'-end formation. *Proc. Natl. Acad. Sci. USA* 107, 17945–17950.
- Field, Y., Kaplan, N., Fondufe-Mittendorf, Y., Moore, I.K., Sharon, E., Lubling, Y., Widom, J., and Segal, E. (2008). Distinct modes of regulation by chromatin encoded through nucleosome positioning signals. *PLoS Comput. Biol.* 4, e1000216. <http://dx.doi.org/10.1371/journal.pcbi.1000216>.
- Gkikopoulos, T., Schofield, P., Singh, V., Pinskaya, M., Mellor, J., Smolle, M., Workman, J.L., Barton, G.J., and Owen-Hughes, T. (2011). A role for Snf2-related nucleosome-spacing enzymes in genome-wide nucleosome organization. *Science* 333, 1758–1760.
- Heus, J.J., Zonneveld, B.J., Bloom, K.S., de Steensma, H.Y., and van den Berg, J.A. (1993). The nucleosome repeat length of *Kluyveromyces lactis* is 16 bp longer than that of *Saccharomyces cerevisiae*. *Nucleic Acids Res.* 21, 2247–2248.
- Jiang, C., and Pugh, B.F. (2009). Nucleosome positioning and gene regulation: advances through genomics. *Nat. Rev. Genet.* 10, 161–172.
- Kaplan, N., Moore, I.K., Fondufe-Mittendorf, Y., Gossett, A.J., Tillo, D., Field, Y., LeProust, E.M., Hughes, T.R., Lieb, J.D., Widom, J., and Segal, E. (2009). The DNA-encoded nucleosome organization of a eukaryotic genome. *Nature* 458, 362–366.
- Kaplan, N., Moore, I., Fondufe-Mittendorf, Y., Gossett, A.J., Tillo, D., Field, Y., Hughes, T.R., Lieb, J.D., Widom, J., and Segal, E. (2010). Nucleosome sequence preferences influence in vivo nucleosome organization. *Nat. Struct. Mol. Biol.* 17, 918–920, author reply 920–922.
- Korber, P., and Horz, W. (2004). In vitro assembly of the characteristic chromatin organization at the yeast PHO5 promoter by a replication-independent extract system. *J. Biol. Chem.* 279, 35113–35120.

- Kornberg, R.D., and Stryer, L. (1988). Statistical distributions of nucleosomes: nonrandom locations by a stochastic mechanism. *Nucleic Acids Res.* **16**, 6677–6690.
- Mavrich, T.N., Ioshikhes, I.P., Venters, B.J., Jiang, C., Tomsho, L.P., Qi, J., Schuster, S.C., Albert, I., and Pugh, B.F. (2008). A barrier nucleosome model for statistical positioning of nucleosomes throughout the yeast genome. *Genome Res.* **18**, 1073–1083.
- Radman-Livaja, M., and Rando, O.J. (2010). Nucleosome positioning: how is it established, and why does it matter? *Dev. Biol.* **339**, 258–266.
- Riley, J., Butler, R., Ogilvie, D., Finniear, R., Jenner, D., Powell, S., Anand, R., Smith, J.C., and Markham, A.F. (1990). A novel, rapid method for the isolation of terminal sequences from yeast artificial chromosome (YAC) clones. *Nucleic Acids Res.* **18**, 2887–2890.
- Segal, E., Fondufe-Mittendorf, Y., Chen, L., Thastrom, A., Field, Y., Moore, I.K., Wang, J.P., and Widom, J. (2006). A genomic code for nucleosome positioning. *Nature* **442**, 772–778.
- Sekinger, E.A., Moqtaderi, Z., and Struhl, K. (2005). Intrinsic histone-DNA interactions and low nucleosome density are important for preferential accessibility of promoter regions in yeast. *Mol. Cell* **18**, 735–748.
- Shivaswamy, S., Bhinge, A., Zhao, Y., Jones, S., Hirst, M., and Iyer, V.R. (2008). Dynamic remodeling of individual nucleosomes across a eukaryotic genome in response to transcriptional perturbation. *PLoS Biol.* **6**, e65. <http://dx.doi.org/10.1371/journal.pbio.0060065>.
- Stein, A., Takasuka, T.E., and Collings, C.K. (2009). Are nucleosome positions in vivo primarily determined by histone-DNA sequence preferences? *Nucleic Acids Res.* **38**, 709–719.
- Tirosh, I., Berman, J., and Barkai, N. (2007). The pattern and evolution of yeast promoter bendability. *Trends Genet.* **23**, 318–321.
- Tirosh, I., Sigal, N., and Barkai, N. (2010). Divergence of nucleosome positioning between two closely related yeast species: genetic basis and functional consequences. *Mol. Syst. Biol.* **6**, 365.
- Tsankov, A.M., Thompson, D.A., Socha, A., Regev, A., and Rando, O.J. (2010). The role of nucleosome positioning in the evolution of gene regulation. *PLoS Biol.* **8**, e1000414. <http://dx.doi.org/10.1371/journal.pbio.1000414>.
- Tsankov, A., Yanagisawa, Y., Rhind, N., Regev, A., and Rando, O.J. (2011). Evolutionary divergence of intrinsic and trans-regulated nucleosome positioning sequences reveals plastic rules for chromatin organization. *Genome Res.* **21**, 1851–1862.
- Vaillant, C., Palmeira, L., Chevereau, G., Audit, B., d'Aubenton-Carafa, Y., Thermes, C., and Arneodo, A. (2010). A novel strategy of transcription regulation by intragenic nucleosome ordering. *Genome Res.* **20**, 59–67.
- Valouev, A., Johnson, S.M., Boyd, S.D., Smith, C.L., Fire, A.Z., and Sidow, A. (2011). Determinants of nucleosome organization in primary human cells. *Nature* **474**, 516–520.
- Van Holde, K.E. (1989). *Chromatin* (New York: Springer-Verlag).
- Weiner, A., Hughes, A., Yassour, M., Rando, O.J., and Friedland, N. (2010). High-resolution nucleosome mapping reveals transcription-dependent promoter packaging. *Genome Res.* **20**, 90–100.
- Whitehouse, I., and Tsukiyama, T. (2006). Antagonistic forces that position nucleosomes in vivo. *Nat. Struct. Mol. Biol.* **13**, 633–640.
- Whitehouse, I., Rando, O.J., Delrow, J., and Tsukiyama, T. (2007). Chromatin remodelling at promoters suppresses antisense transcription. *Nature* **450**, 1031–1035.
- Yuan, G.C., Liu, Y.J., Dion, M.F., Slack, M.D., Wu, L.F., Altschuler, S.J., and Rando, O.J. (2005). Genome-scale identification of nucleosome positions in *S. cerevisiae*. *Science* **309**, 626–630.
- Zhang, Y., Moqtaderi, Z., Rattner, B.P., Euskirchen, G., Snyder, M., Kadonaga, J.T., Liu, X.S., and Struhl, K. (2009). Intrinsic histone-DNA interactions are not the major determinant of nucleosome positions in vivo. *Nat. Struct. Mol. Biol.* **16**, 847–852.
- Zhang, Y., Moqtaderi, Z., Rattner, B.P., Euskirchen, G., Snyder, M., Kadonaga, J.T., Liu, X.S., and Struhl, K. (2010). Evidence against a genomic code for nucleosome positioning. *Nat. Struct. Mol. Biol.* **17**, 920–923.
- Zhang, Z., Wippo, C.J., Wal, M., Ward, E., Korber, P., and Pugh, B.F. (2011). A packing mechanism for nucleosome organization reconstituted across a eukaryotic genome. *Science* **332**, 977–980.